

# Algebra Solving Equations With Variables On Both Sides

Algebra Solving Equations With  
Variables On Both Sides

Downloaded from [everybodystreet.com](http://everybodystreet.com)  
by Patrick & French

## INTRODUCTION: THE NEXT STEP IN YOUR ALGEBRA JOURNEY

If you have been studying algebra for a while, you have likely mastered the art of solving simple equations like  $2x + 3 = 11$ . You know how to isolate the variable by performing inverse operations, and you understand the importance of keeping the equation balanced. But what happens when the variable appears on both sides of the equals sign? This is where many students feel a bit of hesitation, but it is also where algebra becomes truly powerful. Mastering **Algebra Solving Equations With Variables On Both Sides** is a pivotal skill that unlocks a deeper understanding of mathematical relationships and prepares you for more complex problem-solving. This guide will walk you through everything you need to know, from the core principles to advanced tips, ensuring you approach these equations with confidence and clarity.

In this comprehensive article, we will explore the "why" behind the process, break down the steps methodically, examine common pitfalls, and look at real-world scenarios where this type of equation appears. Whether you are a student preparing for a test, a parent helping with homework, or someone refreshing their math skills, this guide is designed to be your go-to resource. Let us demystify the process and turn confusion into mastery.

## UNDERSTANDING THE CORE CONCEPT: WHAT DOES "VARIABLES ON BOTH SIDES" MEAN?

At its simplest, an equation with variables on both sides is any mathematical statement where the unknown quantity (usually represented by a letter like  $x$ ,  $y$ , or  $n$ ) appears in terms on both the left and right sides of the equal sign. For example, the equation  $5x + 2 = 3x + 10$  has the variable  $x$  on both the left side ( $5x$ ) and the right side ( $3x$ ). This is different from a basic equation where the variable is only on one side.

Why does this matter? In real life, situations are rarely one-sided. Consider two mobile phone plans. Plan A charges a fixed fee plus a per-minute rate, and Plan B charges a different fixed fee and a different per-minute rate. To find out when the two plans cost the same, you would set up an equation where the total cost of Plan A (which includes your variable) equals the total cost of Plan B (which also includes your variable). This is a perfect example of why **Algebra Solving Equations With Variables On Both Sides** is not just an academic exercise; it is a tool for making informed decisions.

The fundamental principle that makes solving these equations possible is the **Properties of Equality**. Whatever you do to one side of the equation, you must do to the other. This concept of balance is the bedrock of all algebra. When variables appear on both sides, our goal is to strategically use these properties to move all variable terms to one side and all constant terms to the other, simplifying the equation into a familiar form.

## THE STEP-BY-STEP METHOD FOR SOLVING EQUATIONS WITH VARIABLES ON BOTH SIDES

While every equation is unique, a consistent, systematic approach will help you solve any equation involving variables on

both sides. Follow these steps in order, and you will rarely get lost.

## Step 1: Simplify Both Sides of the Equation

Before you start moving terms, ensure that each side of the equation is as simple as possible. Use the distributive property to eliminate parentheses, and combine any like terms on the same side. For example, in the equation  $2(x + 4) + 3x = 5x - 2$ , you would first distribute the 2 to get  $2x + 8 + 3x = 5x - 2$ , and then combine the  $2x$  and  $3x$  on the left to get  $5x + 8 = 5x - 2$ . This step prevents errors that arise from working with messy expressions.

## Step 2: Move the Variable Terms to One Side

This is the most crucial step in **Algebra Solving Equations With Variables On Both Sides**. Choose which side of the equation will hold the variable terms. Often, it is easiest to move the smaller variable term to the other side to avoid working with negative coefficients, but this is a matter of preference. Use inverse operations (addition or subtraction) to eliminate the variable term from one side.

- **Example:** In  $3x + 6 = 2x + 10$ , you can subtract  $2x$  from both sides to get  $x + 6 = 10$ . Alternatively, you could subtract  $3x$  from both sides, giving you  $6 = -x + 10$ , which is still solvable but requires an extra step.

## Step 3: Move the Constant Terms to the Other Side

Once all variable terms are on one side, use inverse operations to move all constant numbers to the opposite side. This isolates the variable. Continuing the example above, from  $x + 6 = 10$ , you would subtract 6 from both sides to get  $x = 4$ .

## Step 4: Isolate the Variable

If the variable has a coefficient attached to it (other than 1), divide both sides of the equation by that coefficient. If the coefficient is negative, remember to divide by a negative number, which will change the sign of the result. For instance, if you have  $-4x = 12$ , you would divide both sides by  $-4$  to get  $x = -3$ .

## Step 5: Check Your Solution

This final step is your safety net. Substitute the value you found for the variable back into the **original equation**. Simplify both sides independently. If they are equal, your solution is correct. If not, retrace your steps to find the error. Checking not only verifies your answer but also reinforces your understanding of the

equation's structure.

**WORKED EXAMPLES: PUTTING THE PROCESS INTO PRACTICE**

Let us apply the steps above to a few diverse examples. Seeing the process in action is one of the best ways to internalize **Algebra Solving Equations With Variables On Both Sides**.

## Example 1: A Classic Linear Equation

Solve for  $x$ :  $7x - 8 = 4x + 7$

- Simplify:** Both sides are already simplified. No parentheses or like terms to combine.
- Move variables:** Subtract  $4x$  from both sides:  $(7x - 4x) - 8 = (4x - 4x) + 7 \rightarrow 3x - 8 = 7$
- Move constants:** Add 8 to both sides:  $3x - 8 + 8 = 7 + 8 \rightarrow 3x = 15$
- Isolate:** Divide both sides by 3:  $3x / 3 = 15 / 3 \rightarrow x = 5$
- Check:**  $7(5) - 8 = 35 - 8 = 27$  and  $4(5) + 7 = 20 + 7 = 27$ . Both sides equal 27. Correct!

## Example 2: An Equation with Parentheses

Solve for  $y$ :  $4(y - 2) = 8y + 12$

- Simplify:** Distribute the 4 on the left:  $4y - 8 = 8y + 12$
- Move variables:** Subtract  $4y$  from both sides:  $4y - 4y - 8 = 8y - 4y + 12 \rightarrow -8 = 4y + 12$
- Move constants:** Subtract 12 from both sides:  $-8 - 12 = 4y + 12 - 12 \rightarrow -20 = 4y$
- Isolate:** Divide both sides by 4:  $-20 / 4 = 4y / 4 \rightarrow y = -5$
- Check:**  $4(-5 - 2) = 4(-7) = -28$  and  $8(-5) + 12 = -40 + 12 = -28$ . Correct!

## Example 3: Variables on Both Sides with Fractions

Solve for  $a$ :  $(1/2)a + 3 = (1/4)a + 6$

- Simplify:** No parentheses. Note the fractions.
- Move variables:** Subtract  $(1/4)a$  from both sides:  $(1/2 - 1/4)a + 3 = 6 \rightarrow (1/4)a + 3 = 6$
- Move constants:** Subtract 3 from both sides:  $(1/4)a = 3$
- Isolate:** Multiply both sides by 4:  $4 * (1/4)a = 4 * 3 \rightarrow a = 12$
- Check:**  $(1/2)(12) + 3 = 6 + 3 = 9$  and  $(1/4)(12) + 6 = 3 + 6 = 9$ . Correct!

**SPECIAL CASES: WHEN THE VARIABLE DISAPPEARS**

Not all equations with variables on both sides will yield a single neat solution. Sometimes, the variable terms cancel out entirely. Understanding these special cases is a sign of advanced proficiency in **Algebra Solving Equations With Variables On Both Sides**.

### Case 1: No Solution

If, after simplifying and moving terms, you are left with a false statement (like  $5 = 8$ ), then the original equation has no solution. This means there is no value for the variable that will make the equation true.

**Example:**  $2x + 5 = 2x + 9$ . Subtract  $2x$  from both sides:  $5 = 9$ . This is false. Therefore, no solution exists. The symbol for this is often  $\emptyset$ .

### Case 2: Infinite Solutions (Identity)

If the variable terms cancel out and you are left with a true statement (like  $3 = 3$ ), then the equation is an identity. It is true for all real numbers. Every real number is a solution.

**Example:**  $3(x - 1) = 3x - 3$ . Distribute on the left:  $3x - 3 = 3x - 3$ . Subtract  $3x$  from both sides:  $-3 = -3$ . This is true. Therefore, the solution is "all real numbers."

**COMMON MISTAKES AND HOW TO AVOID THEM**

Even experienced students can stumble when solving these equations. Being aware of common errors is half the battle. Here are the most frequent pitfalls in **Algebra Solving Equations With Variables On Both Sides** and how to sidestep them.

- Mistake 1: Forgetting to Distribute a Negative Sign.**  
When you have an expression like