

How To Make Your Period Come

How To Make Your Period Come

Downloaded from everybodystreet.com by Morrow & Mila

UNDERSTANDING THE DESIRE TO PAUSE YOUR MENSTRUAL CYCLE

For many people, their monthly period is inconvenient, painful, or disruptive. The idea of how to make your period stop—temporarily or permanently—is a common question. Athletes might want to avoid bleeding during a competition, travelers may wish to skip their cycle for a vacation, and individuals with severe endometriosis, heavy bleeding, or severe menstrual cramps often seek relief. Thanks to modern medicine, there are several safe, effective ways to suppress menstruation. However, it is crucial to understand that while stopping your period is possible, it should always be done under the guidance of a healthcare professional. This article will explore the most reliable medical methods, lifestyle approaches, and important safety considerations related to menstrual suppression.

Before diving into the methods, it is helpful to understand why you might want to explore how to make your period stop. Menstrual suppression can improve quality of life for those with conditions like **menorrhagia** (heavy bleeding), **dysmenorrhea** (painful periods), or **premenstrual dysphoric disorder (PMDD)**. Others simply prefer not to have a monthly bleed. Regardless of the reason, the decision should be informed by reliable information and a conversation with your doctor, as some methods are more suitable than others depending on your health history and goals.

MEDICAL METHODS FOR STOPPING YOUR PERIOD

When considering how to make your period stop, medical interventions are the most proven and reliable. These methods involve altering your hormone levels or the uterine lining to prevent bleeding. The approaches below are widely used and generally considered safe for long-term use for most people with a uterus.

Continuous or Extended-Cycle Hormonal Birth Control

The most common medical approach involves using hormonal contraception without the usual placebo or pill-free break. This includes birth control pills, the contraceptive patch, or the vaginal ring. Instead of taking a week of placebo pills (or removing the ring/ patch for a week), you simply continue taking active hormones every day. This artificially mimics a pregnancy-like state, keeping the uterine lining thin and preventing a withdrawal bleed.

- **How it works:** The progestin component of the pill (or other hormonal method) prevents the endometrium (uterine lining) from building up. Without a thick lining, there is no bleeding when you skip the break.
- **Effectiveness:** Most people who use continuous birth control pills experience no bleeding or only occasional spotting (breakthrough bleeding) within a few months. For many, periods stop completely.
- **Products:** There are dedicated extended-cycle pills like Seasonique (84 active pills, then 7 low-dose estrogen pills), but you can also use any monophasic birth control pill continuously. Your doctor can prescribe a year’s supply with instructions to skip the placebos.
- **Considerations:** Spotting is common in the first few months. The risk of blood clots is slightly higher with estrogen-containing methods, especially for smokers over 35, those with high blood pressure, or a history of migraines with aura. Progestin-only methods (like the mini-pill, implant, or shot) are often better for those who cannot take estrogen.

Progestin-Only Contraceptives

For many, the answer to how to make your period stop lies in progestin-only methods. These are safer for people with contraindications to estrogen, such as a history of blood clots or migraine with aura.

- **The Levonorgestrel Intrauterine Device (IUD):** Hormonal IUDs such as Mirena, Liletta, Kyleena, and Skyla release a small amount of progestin directly into the uterus. This thins the uterine lining significantly. Within one year of insertion, about 50-80% of users experience light or no periods. After several years, many stop bleeding altogether. It is a highly effective, long-acting reversible contraceptive (LARC) that lasts 3-7 years depending on the brand.
- **The Contraceptive Implant (Nexplanon):** A small rod placed under the skin of the upper arm releases progestin. It is very effective at preventing pregnancy and often leads to lighter or absent periods, though irregular bleeding is a common side effect for some.
- **The Birth Control Shot (Depo-Provera):** An injection of progestin given every 12 weeks. Over time, many users stop menstruating entirely. However, it can cause irregular spotting initially, and long-term use (over 2 years) may lead to a temporary reduction in bone density, which typically recovers after stopping.
- **The Progestin-Only Pill (Mini-Pill):** Must be taken at the same time every day without a break. It can lighten periods or stop them, but it is less consistent for complete suppression than the IUD or implant.

Gonadotropin-Releasing Hormone (GnRH) Agonists

These medications temporarily place the body into a reversible menopausal state by suppressing ovarian hormone production. They are typically used for a few months to treat conditions like severe endometriosis or to reduce bleeding before surgery. Examples include Lupron (leuprolide) and Zoladex (goserelin). They are not recommended for long-term use due to side effects like hot flashes, vaginal dryness, and bone loss, but they are an effective short-term solution for how to make your period stop completely. Add-back hormone therapy (small doses of estrogen or progestin) can reduce side effects if used longer.

Non-Hormonal Surgical Options

For those who are certain they do not want future children or are done with childbearing, surgical procedures can permanently stop menstruation. These are major decisions and not considered first-line approaches for purely stopping periods.

- **Endometrial Ablation:** This procedure destroys the lining of the uterus, making it unable to thicken or bleed. It is minimally invasive but not a form of contraception (you still need birth control to prevent pregnancy). It works very well for heavy bleeding; many people have no periods afterward. However, it may not be as effective for very young women or those with certain uterine anatomy issues. It is not reversible.
- **Hysterectomy:** Surgical removal of the uterus. This permanently stops menstruation and eliminates the possibility of pregnancy. It is a major operation with recovery time and permanent consequences. It is usually reserved for cases where other treatments have failed for severe conditions like fibroids, adenomyosis, or cancer.

ARE THERE NATURAL WAYS TO STOP YOUR PERIOD?

Many people search for natural remedies to answer how to make your period stop. It is important to set realistic expectations: there are no scientifically proven natural methods that will reliably stop a period once it has started or prevent it from coming in the next cycle. However, certain lifestyle factors may influence cycle regularity and flow.

That said, you should never try to “stop” a period that is already happening using unproven methods like large doses of ibuprofen (though NSAIDs can reduce flow, they won’t stop it completely) or consuming large amounts of lemon juice, gelatin, or vinegar, as these are not safe or effective. Always consult a doctor before attempting any herbal or dietary experiment.

Lifestyle and Dietary Factors That May Influence Flow

- **Non-Steroidal Anti-Inflammatory Drugs (NSAIDs):** Taking high doses of ibuprofen or naproxen during your period can reduce prostaglandins, which are chemicals that cause cramps and heavy bleeding. This can reduce the amount of menstrual flow by about 20-30% in some individuals, but it will not stop it.
- **Stress Management:** Chronic stress can disrupt the hypothalamic-pituitary-ovarian axis, potentially delaying or skipping a period entirely. However, this is unreliable and unpredictable. Purposefully increasing stress is not a recommended strategy.
- **Weight and Exercise:** Very low body fat from extreme exercise or eating disorders can cause amenorrhea (absence of periods). However, deliberately underweighting yourself is dangerous and not a valid method for how to make your period stop. Maintaining a healthy weight supports regular cycles.
- **Herbs:** Some herbs like chasteberry (Vitex) or ginger are touted, but there is no robust evidence that they stop menstruation. They may interact with hormones and medications, so caution is needed.

The honest truth is that while lifestyle modifications can affect menstrual health, they are not reliable means of suppressing menstruation. Medical methods are the only proven, safe way to achieve consistent and predictable absence of periods.

IS IT SAFE TO STOP YOUR PERIOD?

This is a critical question when researching how to make your period stop. Many people worry that stopping a period is unnatural or risky. However, for most individuals, it is perfectly safe. Menstruation is not a natural “detox” process; the body does not need to bleed to be healthy. The uterine lining is shed because there is no pregnancy, but allowing the lining to remain thin through hormonal methods does not cause harm.

Research indicates that continuous birth control use is safe for many years. The main risks are associated with the method itself (e.g., blood clots from estrogen, bone density concerns from Depo-Provera), not the act of not bleeding. Breakthrough bleeding can be annoying but is generally not dangerous, though it should be reported to a doctor to rule out other causes.

Certain medical conditions may make stopping your period more complicated. For instance, if you have a bleeding disorder, you may need special monitoring. Always discuss your desire to suppress menstruation with a gynecologist or primary care provider who knows your full health history.

WHEN TO CONSULT A DOCTOR

You should schedule a medical appointment to discuss how to make your period stop if:

- You experience heavy bleeding (soaking through a pad or tampon every hour for several hours) or severe pain that interferes with daily life.
- You have a personal or family history of blood clots, stroke, or heart disease.
- You smoke and are over 35 years old.
- You have migraines with aura (this affects whether estrogen is safe for you).
- You are trying to conceive or want to preserve fertility.
- You have breast cancer or liver disease.
- You have unexplained irregular bleeding between periods or after intercourse.

Your doctor can help you choose the best method based on your health, lifestyle, and future pregnancy plans. They can also manage any side effects you may experience.

MANAGING SIDE EFFECTS AND BREAKTHROUGH BLEEDING

If you are using a continuous hormonal method and experience spotting or light bleeding, do not panic. Breakthrough bleeding is very common during the first 3-6 months. To manage it:

- Take your active pills at exactly the same time daily (especially important for progestin-only pills).
- If you are on a combination pill and spotting persists, talk to your doctor about switching to a

pill with a different progestin or a higher dose of estrogen.

- Allow yourself a 3- or 4-day break after several months of continuous use if the spotting is bothersome (this can be done under medical guidance to trigger a scheduled withdrawal bleed and reset the lining).
- Use panty liners and keep a small bag with supplies during the adjustment period.

For those with an IUD, spotting is common for the first 3-6 months but typically resolves. The

implant may cause unpredictable bleeding patterns, but this often becomes lighter over time.

MYTHS AND MISCONCEPTIONS ABOUT STOPPING YOUR PERIOD

There are many myths surrounding how to make your period stop. Let’s address some common ones:

- **Myth:** Stopping your period causes toxins to