
How To Help Someone With Borderline Personality Disorder

*How To Help Someone
With Borderline
Personality Disorder*

*Downloaded from
everybodystreet.com by
Bronson & Kendrick*

UNDERSTANDING THE CHALLENGES AND OFFERING REAL SUPPORT

Loving or caring for someone diagnosed with borderline personality disorder (BPD) can feel like navigating an emotional storm without a map. The intense mood swings, fear of abandonment, and impulsive actions can

leave family members and friends feeling confused, exhausted, and even hopeless. However, understanding how to help someone with borderline personality disorder is not only possible but can be profoundly transformative for both of you. With the right knowledge, strategies, and a commitment to compassionate boundaries, you can become a stabilizing force in their life without sacrificing your own well-being.

Before diving into specific techniques, it

is crucial to recognize that BPD is a legitimate mental health condition, not a character flaw or a choice. Individuals with BPD often experience a chronic sense of emptiness, unstable relationships, and a deeply fractured sense of self. Their reactions, which may seem manipulative or overly dramatic to outsiders, are usually desperate attempts to manage overwhelming internal pain. By reframing their behavior through this lens of suffering, you can begin to approach the situation with genuine empathy rather than frustration.

WHAT IS BORDERLINE PERSONALITY DISORDER? A BRIEF OVERVIEW

To help effectively, you must first

understand what you are dealing with. Borderline personality disorder is characterized by a pervasive pattern of instability in interpersonal relationships, self-image, and emotions. The Diagnostic and Statistical Manual of Mental Disorders (DSM-5) outlines nine specific criteria, and a diagnosis typically requires meeting at least five of them. These include frantic efforts to avoid real or imagined abandonment, a pattern of intense and unstable relationships, marked impulsivity, recurrent suicidal behavior or self-harm, affective instability, chronic feelings of emptiness, and difficulty controlling anger.

It is also important to note that BPD often co-occurs with other conditions such as depression, anxiety disorders, eating disorders, or substance use

disorders. This complexity means that a holistic approach to treatment and support is necessary. The good news is that with proper therapy — particularly dialectical behavior therapy (DBT), mentalization-based therapy (MBT), or transference-focused psychotherapy (TFP) — many individuals with BPD experience significant improvement over time.

EDUCATE YOURSELF WITHOUT OVERWHELMING THEM

One of the most powerful first steps in learning how to help someone with borderline personality disorder is to educate yourself about the condition. Read books by experts like Dr. Marsha Linehan, who developed DBT, or explore reputable online resources from

organizations such as the National Education Alliance for Borderline Personality Disorder (NEA-BPD). Knowledge reduces fear and helps you respond with insight rather than reactivity.

However, be careful not to become an amateur diagnostician. Avoid labeling their every mood swing or disagreement as a symptom of BPD. Instead, use your understanding to foster patience and compassion. When you recognize that their fear of abandonment is driven by deep-seated insecurity rather than a lack of trust in you, you can respond with reassurance rather than defensiveness.

MASTER THE ART OF VALIDATION

Validation is arguably the single most effective communication tool when

supporting someone with BPD. Validation does not mean agreeing with their perception of reality or condoning harmful behavior. Instead, it means acknowledging their emotional experience as real and understandable given their history and internal state.

For example, if your loved one says, "You are going to leave me just like everyone else," a validating response might be, "I can hear how terrified you are that I will abandon you. That feeling must be incredibly painful. I want you to know that I am here right now, and I care about you." Notice how this response does not argue, defend, or prove anything. It simply meets their emotional reality with acceptance.

Validation helps de-escalate intense emotions because it signals to the

person with BPD that they are seen and heard. When they feel understood, the need to prove their pain through more dramatic actions often diminishes.

Practical Validation Techniques

- **Pay attention:** Give them your full presence without distractions. Put down your phone, make eye contact, and listen without interrupting.
- **Reflect their words:** Paraphrase what they have said to show you are listening. For instance, "So you

felt humiliated when your friend did not invite you."

- **Read their mind:** Guess what they might be feeling even if they have not explicitly stated it. "It looks like you are feeling really hurt right now."
- **Understand the context:** Acknowledge how their past experiences might make the current situation more painful. "Given how your last relationship ended, it makes sense that you would feel anxious right now."
- **Normalize the emotion:** Let them know that their feelings are human and understandable. "Anyone in your situation would feel frustrated."

SET AND MAINTAIN HEALTHY BOUNDARIES

Many people who want to know how to help someone with borderline personality disorder struggle with the concept of boundaries. There is a fear that setting limits will be perceived as rejection, triggering an intense reaction. In reality, clear and consistent boundaries are an essential form of love and support. They provide structure and predictability, which can be deeply calming for someone whose internal world feels chaotic.

Boundaries are not about controlling the other person; they are about taking responsibility for your own well-being and clearly communicating what behavior you will and will not accept. For

example, you might say, "I am happy to talk with you about your feelings, but I will need to end the conversation if you start yelling at me. We can continue later when we are both calm."

Guidelines for Boundary Setting

1. **Be clear and specific:** Vague boundaries are hard to enforce. Instead of saying "Stop being so needy," say "I need some quiet time after work. I will be available to talk after dinner."
2. **Be consistent:** Enforce your boundaries every time. Inconsistency creates confusion

and can actually increase testing behaviors.

3. **Use a calm tone:** Deliver your boundary with a neutral and compassionate voice. Your goal is to protect the relationship, not to punish.
4. **Follow through:** If you say you will end a conversation if yelling continues, then do it. Empty threats undermine trust.

ENCOURAGE PROFESSIONAL TREATMENT WITHOUT NAGGING

You cannot be your loved one's sole therapist or rescuer. While your support is invaluable, professional treatment is the cornerstone of recovery from BPD. Learning how to help someone with borderline personality disorder includes

knowing when to step back and encourage expert care.

Dialectical behavior therapy (DBT) is considered the gold standard treatment for BPD. It focuses on four key skill areas: mindfulness, distress tolerance, emotion regulation, and interpersonal effectiveness. If your loved one is in therapy, you can ask how you might support their skill-building at home. Some therapists even offer family sessions or skills training for loved ones.

If they are not yet in treatment, approach the subject gently. Avoid ultimatums or shaming. You might say, "I have noticed you have been suffering a lot lately, and I care about you. I have read that therapy can be really helpful for managing some of these feelings. Would you be open to exploring that

possibility together?" Offer to help research therapists or accompany them to an initial appointment if they are anxious.

NAVIGATE CRISES WITH A PLAN

Crises, including self-harm or suicidal ideation, are not uncommon in BPD. Knowing how to respond in these moments is critical. Your role during a crisis is not to solve everything but to help stabilize the situation and connect them with appropriate resources.

First, stay as calm as possible. Your calm energy can be contagious and help de-escalate their intensity. Listen without judgment, validate their pain, and avoid minimizing their distress by saying things like "You have so much to live for." Instead, focus on the present

moment and on getting through the next hour.

Ask directly about safety. Contrary to popular belief, asking about suicidal thoughts does not plant the idea. You can say, "Are you thinking about hurting yourself?" If the answer is yes, do not leave them alone. Remove any means of self-harm, and call a crisis hotline or take them to the emergency room if necessary. The 988 Suicide and Crisis Lifeline in the United States is an excellent resource for both of you.

What to Do in a

Crisis

- Stay with the person or ensure they are not alone.
- Use a calm, steady tone of voice.
- Avoid arguing, judging, or giving simple solutions.
- Validate their pain: "I can see you are in tremendous pain right now."
- Encourage them to use coping skills they have learned in therapy.
- Call a crisis line for guidance if you are unsure.

TAKE CARE OF YOURSELF: THE OFTEN-FORGOTTEN PIECE

You cannot pour from an empty cup. If you exhaust yourself trying to support someone with BPD, you will eventually

become resentful, burned out, or even ill. Self-care is not selfish; it is strategic and necessary for long-term support.

Consider joining a support group for family members of people with BPD. Organizations like the National Alliance on Mental Illness (NAMI) offer family-to-family programs. Talking with others who truly understand the challenges can be immensely relieving. Additionally, consider seeing a therapist for yourself. Having a neutral professional to help you process your own feelings and develop coping strategies is a sign of strength, not weakness.

Make time for activities that replenish you. This could be exercise, meditation, time in nature, hobbies, or simply spending time with friends who are not part of the caregiving dynamic. Give

yourself permission to take breaks without guilt. It is okay to say, "I need some time for myself today, but I will check in with you tomorrow."

RECOGNIZE PROGRESS AND CELEBRATE SMALL WINS

Recovery from BPD is rarely a straight line. There will be good days and bad days, steps forward and steps back. It is easy to focus on the setbacks, but it is equally important to recognize and celebrate progress, no matter how small. Did your loved one use a coping skill instead of self-harming? Did they take responsibility for something without blaming someone else? Did they ask for support in a calm, direct way? These are victories worth acknowledging.

Positive reinforcement can be incredibly

motivating. A simple statement like, "I saw how hard you worked to stay calm during that conversation, and I am really proud of you," can strengthen their commitment to growth and deepen your connection.

COMMON MISTAKES TO AVOID

Even with the best intentions, it is easy to fall into patterns that are unhelpful. Being aware of these common pitfalls can help you stay on track.

- **Taking things personally:** When someone with BPD lashes out, it is usually a reflection of their internal pain, not a true judgment of your character. Try to build a mental buffer between their words and your self-worth.
- **Rescuing or enabling:** Solving their problems for them may provide short-term relief but robs them of the opportunity to build their own coping skills.
- **Walking on eggshells:** Constantly tiptoeing around their emotions to avoid conflict is exhausting and ultimately unsustainable. Authenticity is more helpful than perfect peace.
- **Expecting rapid change:** Lasting change takes time, often years of dedicated therapy. Patience is essential.
- **Neglecting your own needs:** Sacrificing your own mental health will not save theirs. It will only lead to two people in crisis.

WHEN TO STEP BACK FOR SAFETY

There may come a point when the relationship becomes unsafe, whether emotionally, physically, or both. If you are experiencing verbal abuse, threats, physical aggression, or extreme manipulation that is destroying your mental health, it is okay to step back. Loving someone from a distance is sometimes the most compassionate choice you can make.

This does not mean you are giving up on them. It means you are recognizing the limits of what you can provide. You can still wish them well, encourage treatment from afar, and hope for their recovery without being an active participant in a toxic dynamic. Your

safety matters.

CONCLUSION

Learning how to help someone with borderline personality disorder is a journey that requires patience, education, empathy, and clear boundaries. You cannot fix them or cure them, but you can offer a steady, validating presence that makes their path to healing a little less lonely. By focusing on validation, encouraging professional treatment, managing crises calmly, and prioritizing your own self-care, you create an environment where both you and your loved one can grow. Progress may be slow and imperfect, but every moment of genuine connection is a step